

Effect of ration size and meal frequency on assimilation and digestive efficiency in yearling Steller sea lions, *Eumetopias jubatus*

David A. S. Rosen, Laura Williams and Andrew W. Trites

Marine Mammal Research Unit, University of British Columbia, 6248 Biological Sciences Rd., Vancouver, British Columbia V6T 1Z4, Canada

Abstract

Assimilation and digestive efficiencies were measured in four juvenile Steller sea lions (*Eumetopias jubatus*) fed three ration sizes of herring (3%, 6%, or 9% of body mass) at three frequencies (2, 3, or 4 times daily). Assimilation efficiency (dry matter digestive efficiency) was $90.0 \pm 2.0\%$ (mean \pm 1 SD). Digestive efficiency (efficiency of energy digestion) was $95.5 \pm 1.0\%$. There was a strong linear relationship between digestive and assimilation efficiency, but no significant differences in either assimilation or digestive efficiency with changes in feeding frequency or changes in daily food intake within the ranges offered.

Key words: Steller sea lion, *Eumetopias jubatus*, digestion, prey, herring.

Introduction

Feeding studies are important in understanding the ecology of Steller sea lions (*Eumetopias jubatus*) given that nutritional stress may be the most significant contributing factor to the population declines observed in Alaska since the 1970's (Merrick *et al.*, 1987; Trites & Larkin, 1996). Changes in the nutritional status of Steller sea lions may be attributable to changes in their prey base (distribution, composition, and abundance). An understanding of the bioenergetic consequences of changes in diet is relevant to both sea lions and other top marine predators in the north Pacific ecosystem, many of which also exhibit population declines (Pitcher, 1990; Trites, 1992; Byrd & Dragoo, 1997).

Accurate bioenergetic models are needed to determine how changes in feeding ecology affect the energy budgets of Steller sea lions and other species. A critical component of energetic models is the calculation of metabolizable energy. This requires estimates of the amount of energy lost through urine, feces, and the heat increment of feeding.

These parameters vary with the type and composition of the prey consumed, meal size, frequency of eating, the morphology of the digestive tract, and the age and sex of the animal (Keiver *et al.*, 1984; Ronald *et al.*, 1984; Fisher *et al.*, 1992; Mårtensson *et al.*, 1994a; Mårtensson *et al.*, 1994b; Lawson *et al.*, 1997a; Lawson *et al.*, 1997b).

Subtracting fecal energy loss (FE) from gross energy intake (GEI) yields apparent digestible energy (Lavigne *et al.*, 1982). Digestible energy, when expressed in relation to gross energy intake, is called digestive efficiency (DE):

$$DE = (GEI - FE) / GEI * 100.$$

Digestive efficiency studies can use either the differences in energy content between the food and whole fecal collection, or comparisons of the changes in energy and concentration of an inert dietary marker between the food item and a fecal sample (Kleiber, 1975), such as naturally occurring manganese (Fadely *et al.*, 1990). Digestive efficiency is used to calculate the apparent digestible energy derived from food intake, and is a critical parameter in the construction of energy budgets.

Some studies use dry matter assimilation efficiency (AE) as a measure of the digestive value derived from particular prey. This measure is easier to determine, being calculated solely from changes in an inert marker between fecal and food samples. Assimilation efficiency may provide a reasonable estimate of digestive efficiency, if the proportion of non-digestible parts in the prey is low.

As passage rates are affected by meal frequency and ration size (Hunt & Stubbs, 1975; Warner, 1981; Markussen, 1993), these variables might affect the digestive and assimilation efficiencies of marine mammals (Ronald *et al.*, 1984; Lawson *et al.*, 1997b). On the other hand, symmorphosis suggests that the digestive system should be able to accommodate reasonable variation in foraging frequency and meal size (Karasov & Diamond, 1988; Diamond, 1991). This question is of particular

relevance to Steller sea lions in the North Pacific who may be forced to alter their foraging patterns in response to changes in their prey base. This study quantified the assimilation and digestive efficiencies for juvenile Steller sea lions fed herring, and tested whether digestive and assimilation efficiencies were affected by meal frequency and daily food intake. The relationship between assimilation and digestive efficiency was also examined.

Materials and Methods

The experiment was conducted between February and March 1998 on four captive Steller sea lions, approximately 8 months of age, kept at the Vancouver Aquarium. The pups, two males and two females, were weaned from milk formula to herring (*Clupea harengus pallasi*) in September 1997. Their normal diet (September to January) consisted of thawed herring fed 3 times per day, totaling approximately 6% of their body weight. The sea lions were weighed every morning before the first feed on a platform scale (± 0.5 kg).

The experiment consisted of 2 feeding manipulations: total food intake and number of feedings per day. Each sea lion received a total daily food intake of herring equal to either 3%, 6%, or 9% of their body weight (body mass of 4 pups: 46.8–70.7 kg). This quantity was delivered to the animals over the course of either 2, 3, or 4 feedings per day. Thus, each pup undertook 9 (randomized) trial combinations, with each trial lasting 4 days. Four day trials were estimated to be long enough to ensure that the sea lions had adjusted to each new feeding regime given their very fast rates of digestion (Helm, 1984).

Feces was collected from the sea lions on days 3 and 4 of each trial. Sea lions were placed alone in a cement floored run to ensure the scat came from a known animal. Fecal samples were collected from noon on the third day onward and, if no samples were collected during the day, the sea lion was kept in isolation over night so a sample could be collected early the next morning. If there was no feces, or if the sea lion decided not to eat, the trial was rerun at a later date. All samples were placed in a labeled, plastic vial, and stored at -20°C until processed. Samples of herring were also collected and frozen throughout the trials.

For analysis, fecal samples were thawed overnight. Two duplicate sub-samples (~ 10 g) were placed in aluminum dishes and weighed to the nearest mg (Sartorius laboratory scale, model BP110). Feces were dried to a constant weight in a convection oven for 20–22 h at 100°C to determine water content. After drying, fecal samples were re-weighed and ground with a glass mortar and pestle into a fine, uniform powder. The powdered

Table 1. Mean assimilation efficiencies for 4 sea lions subjected to 9 combinations of trial conditions (meals $2 \times$, $3 \times$, $4 \times$ per day; total food equals 3%, 6%, or 9% of body mass). One standard deviation given in brackets.

Total food (% body mass)	Meals per day			Total
	$2 \times$	$3 \times$	$4 \times$	
3%	89.1 (2.0)	89.5 (1.8)	89.1 (0.8)	89.3 (1.5)
6%	90.6 (0.6)	88.7 (4.4)	90.7 (2.0)	90.0 (2.7)
9%	90.7 (1.2)	90.4 (2.2)	91.0 (1.0)	90.7 (1.4)
Total	90.2 (1.5)	89.5 (2.8)	90.3 (1.5)	90.0 (2.0)

sample was then divided into two new samples of >1 g each, which were sent to outside laboratories to determine energy content and manganese (Mn^{2+}) concentration. Extra samples were stored for possible reanalysis.

Herring samples were partially thawed, cut, and blended into a uniform paste with a food processor. The paste from each fish was divided into 2 sub-samples and the same drying, weighing, grinding procedures, and analyses used as for the scat.

Energy content was determined via an adiabatic bomb calorimetry (Dr. G. Galzi, Department of Animal Science, University of British Columbia). Estimated energy content for each sample was corrected for side reactions (e.g., fuse wire correction and acid titration).

Manganese analysis was carried out by Norwest Laboratory in Surrey, British Columbia. A quantity of the powdered scat sample was digested with HNO_3 and H_2O_2 in a sealed Teflon vessel using microwave heating. The Mn^{2+} concentration was determined on the resulting solution by UNICP-AES.

Assimilation efficiency was calculated as the change in concentration of the manganese marker between the fish and fecal samples:

$$\text{AE} = (1 - (C_f/C_i)) \times 100 \quad (1)$$

where, C is the concentration of Mn^{2+} in the ingested food (i) and feces (f) on a dry matter basis (Fadely *et al.*, 1990).

Digestion efficiency was calculated by comparing both the change in manganese concentration and energy content. We used a corrected version of the formula given in Mårtensson *et al.* (1994a):

$$\text{DE} = \left(1 - \frac{C_i \times E_f}{C_f \times E_i}\right) \times 100 \quad (2)$$

where, C is the concentration of Mn^{2+} and E is the energy content of ingested food (i) and feces (f) on a dry matter basis.




Figure 1. Changes in assimilation efficiency (top) and digestive efficiency (bottom) with total food intake. Bars represent meal frequencies (± 1 SD) of 2, 3, and 4 times per day.

Percentage data were normalized using an arcsine transformation. Data were collapsed across subsamples to yield a single mean estimate for each animal in each treatment. A 2-level, within-subjects (repeated measure) factorial design ANOVA was chosen to test for an effect of meal size or frequency of feeding on the assimilation efficiency and digestion efficiency. Results were considered significant at $\alpha=0.05$.

Results

Mean length of herring used in the experiment was 17.5 cm, with a mean water content of 72.6%, a

mean energy density of 25112 kJ/g dry weight, and a mean Mn^{2+} concentration of 8.09 ppm. Sea lion feces had a mean water content of 61.4%, mean energy density of 12365 kJ/g dry weight, and mean Mn^{2+} concentration of 83.86 ppm.

Mean assimilation efficiency for the sea lions across all treatments was $90.0 \pm 2.0\%$ (Table 1), with a range of 82.2–92.6%. However, no significant differences could be attributed to changes in either feeding frequency ($F_{2,6}=0.19$, $P=0.83$) or daily food intake ($F_{2,6}=4.24$, $P=0.07$) (Fig. 1).

Digestion efficiency averaged $95.5 \pm 1.0\%$ (Table 2), with a range of 93.5–97.3%. Again, there were

Table 2. Mean digestive efficiencies for 4 sea lions subjected to 9 combinations of trial conditions (meals 2 ×, 3 ×, 4 × per day; total food equals 3%, 6%, or 9% of body mass). One standard deviation given in brackets.

Total food (% body mass)	Meals per day			Total
	2 ×	3 ×	4 ×	
3%	94.8 (1.2)	94.9 (0.9)	94.8 (0.7)	94.8 (0.9)
6%	95.7 (0.6)	95.8 (1.0)	96.3 (1.1)	95.9 (0.9)
9%	95.8 (0.7)	95.8 (0.9)	95.9 (1.6)	95.8 (1.0)
Total	95.4 (0.9)	95.5 (1.0)	95.7 (1.3)	95.5 (1.0)

no significant changes in digestive efficiency with either varying daily food intakes ($F_{2,6}=3.99$, $P=0.08$) or different feeding frequencies ($F_{2,6}=0.16$, $P=0.86$) (Fig. 1).

The relationship between assimilation efficiency and digestive efficiency was: $DE=34.63+0.675*AE$ ($r^2=0.78$, Fig. 2). This was calculated using a geometric mean (type II) regression (Ricker, 1973) fit to the raw data. Regressing arc-sine transformed data did not improve the fit.

Discussion

The high values found for digestive efficiency (95.5%) and assimilation efficiency (90.0%) for Steller sea lions eating herring was consistent with those reported for carnivorous mammals in general (see Table 3.5 in Blaxter 1989, Table 4 in Lavigne *et al.* 1982), and pinnipeds in particular. These

results are contrary to the historical reputation of pinnipeds as 'inefficient converters of fish flesh' (Sergeant, 1973).

More specifically, the results are consistent with other studies that have examined assimilation efficiency among pinnipeds fed herring. Assimilation efficiency has been estimated for two otariid species at 89% for California sea lions, *Zalophus californianus* (Fadely *et al.*, 1994) and 90–93% for Northern fur seals, *Callorhinus ursinus* (Miller, 1978; Fadely *et al.*, 1990), and from 88–94% across several phocid seal species (Table 3).

Fewer studies have estimated digestive efficiency in pinnipeds, and no values have been reported for otariids. Miller (1978) quantified fecal energy loss in Northern fur seals, but it is impossible to derive a specific value for digestive efficiency from the data presented. Digestive efficiency of various phocids fed herring diets has been estimated between 93–97% and 92.7% in walruses, *Odobenus rosmarus* (Fisher *et al.*, 1992). In comparison, Nordøy *et al.* (1993) estimated the digestive efficiency of herring for minke whales, *Balaenoptera acutorostrata*, to be 92.1% through *in vitro* digestion experiments (Table 3).

Although past studies yielded similar estimates for both assimilation and digestive efficiency of herring, it must be noted that the composition and type of prey can substantially affect these values. In general, assimilation and digestive efficiency decreases with decreasing energy or lipid content of the prey, although this is not always the case (see Table 3). Therefore, care should be taken when applying values from specific studies to alternate prey items.




Figure 2. Relationship of assimilation and digestive efficiency. The solid line represents the geometric mean linear regression: $DE=34.63+0.675*AE$.

Table 3. Estimates of assimilation (AE) and digestive efficiency (DE) for various pinnipeds eating different diets. Most studies used changes in marker and energy concentrations to calculate assimilation and/or digestive efficiency. Ranges indicate estimates derived from different methodologies.

Species	Diet	AE	DE	Source
Steller sea lion	Herring	90.0	95.5	Tables 1 and 2
California sea lion	Herring	89.2	—	Fadely <i>et al.</i> , 1994
	Pollock	86.5	—	Fadely <i>et al.</i> , 1994
Northern fur seal	Capelin	88.0	—	Miller, 1978
	Herring	90.0	—	Fadely <i>et al.</i> , 1990
	Herring	91.6–93.0	—	Miller, 1978
	Pollock	86.6–90.0	—	Miller, 1978
Pacific walrus	Squid	92.0	—	Miller, 1978
	Clams	89.9	92.0	Fisher <i>et al.</i> , 1992
	Herring	87.8	92.7	Fisher <i>et al.</i> , 1992
	Krill	—	84	Mårtensson <i>et al.</i> , 1994a
Crabeater seal	Herring	87.6	92.6	Ronald <i>et al.</i> , 1984
	Mixed	—	92.8	Prime & Hammond, 1987
Harbour seal	Herring	92.4	—	Rosen, 1996
	Herring	91.2	—	Ashwell-Erickson & Elsner, 1981
	Pollock	96.7	—	Ashwell-Erickson & Elsner, 1981
Harp seal	Arctic cod	86.9	93.5	Lawson <i>et al.</i> , 1997b
	Atlantic cod	84.3	93.2	Lawson <i>et al.</i> , 1997b
	Capelin	91.4	95.7	Lawson <i>et al.</i> , 1997b
	Capelin	—	93–94	Mårtensson <i>et al.</i> , 1994b
	Crustaceans	—	81–83	Mårtensson <i>et al.</i> , 1994b
	Halibut	88.5	94.7	Lawson <i>et al.</i> , 1997b
	Herring	83.9–90.3	92.5–95.0	Keiver <i>et al.</i> , 1984
Ringed seal	Herring	91.0	96.6	Lawson <i>et al.</i> , 1997b
	Shrimp	60.0	72.2	Keiver <i>et al.</i> , 1984
	Arctic cod	88.3	—	Lawson <i>et al.</i> , 1997a
	Capelin	86.6	—	Lawson <i>et al.</i> , 1997a
	Herring	93.9	—	Lawson <i>et al.</i> , 1997a
	Herring	—	97.0	Parsons, 1977
	Redfish	83.2	—	Lawson <i>et al.</i> , 1997a

In the present study there was a significant, linear relationship between assimilation and digestive efficiency. Not surprisingly, digestive efficiency was always greater than assimilation efficiency. Many of the particles excreted through the feces are indigestible prey parts with extremely low energy densities (e.g., otoliths, bones, chitinous exoskeletons). Their passage decreases assimilation efficiency, but contributes little in the way of fecal energy loss, i.e., has little impact on digestive efficiency. Although assimilation and digestive efficiencies may be strongly related, they clearly are not interchangeable. Unfortunately, the measures are sometimes used inappropriately, particularly in cases where assimilation efficiency is used to estimate fecal energy loss.

The nutritional stress hypothesis proposes that the Steller sea lion population decline is related to changes in their prey base in the North Pacific (Alaska Sea Grant, 1993). Resulting changes in foraging frequency and meal size might affect the metabolizable energy derived from prey. It has been

hypothesized that digestive and assimilation efficiencies might be quickly limited by the absorptive capabilities of the digestive system with increases in food intake (Brody, 1945; Blaxter, 1989). Certainly, there is evidence that passage rates decrease with increasing meal size (Hunt and Stubbs, 1975; Warner, 1981; Markussen, 1993), which might be of particular consequence to pinnipeds with their remarkably long small intestine (Eastman & Coalson, 1974; Martin, 1977), and rapid transit times (Helm, 1984; Markussen, 1993).

In the present study, the Steller sea lions did not exhibit a significant change in their assimilation or digestive efficiency in response to the offered changes in daily food totals or feeding frequency. Similar results were reported for harp (*Phoca groenlandica*), ringed (*P. hispida*), and grey seals (*Halichoerus grypus*) (Keiver *et al.*, 1984; Ronald *et al.*, 1984; Lawson *et al.*, 1997a; Lawson *et al.*, 1997b). While it is possible that these studies did not sufficiently increase food intake to test the

limitation of the system (or that the trials were not of sufficient length), it is more likely that, in accordance with the theory of symmorphosis (Diamond, 1993), there is sufficient capacity within the digestive system to handle any natural load efficiently. The pinniped digestive tract, like that of most other carnivores (Karasov and Diamond, 1988), appears to be adapted to rapid, efficient digestion of bolus meals, which is particularly useful when ingesting large amounts of food at depth (Krockenberger & Bryden, 1994).

This study found that Steller sea lions exhibit digestive and assimilation efficiencies typical of most carnivorous mammals. This high level of efficiency did not seem to be affected by either the frequency or size of meals within the range offered. While these attributes likely reflect the general feeding ecology of Steller sea lions in particular, and pinnipeds in general, comparative data are thin. Further investigations are needed into the possible effects of age, sex, and nutritional status across a larger number of species fed a more complex array of diets.

Acknowledgments

We thank G. Shephard and B. Lasby for their assistance with sample collection, J. Lawson for advice on sample preparation, and A. Winship for comments on the manuscript. This research was funded by the North Pacific Marine Science Foundation through the North Pacific Universities Marine Mammal Research Consortium.

Literature Cited

- Alaska Sea Grant (1993) Is it food? Addressing marine mammal and sea bird declines. University of Alaska Fairbanks, Alaska Sea Grant Report No. 93-01.
- Ashwell-Erickson, S. & Elsner, R. (1981) The energy cost of free existence for Bering Sea harbor and spotted seals. In: D. W. Hood & J. A. Calder (eds.) *The Eastern Bering Sea Shelf: Oceanography and Resources*. Vol. 2, pp. 869–899. University of Washington Press, Seattle.
- Blaxter, K. (1989) *Energy Metabolism in Animals and Man*. Cambridge University Press, Cambridge.
- Brody, S. (1945) *Bioenergetics and Growth*. Hafner Press, New York.
- Byrd, G. V. & Drago, D. E. (1997) Breeding success and population trends of selected seabirds in Alaska in 1996. *U.S. Fish and Wildlife Services Report No. AMNWR 97/11*.
- Diamond, J. (1991) Evolutionary design and intestinal nutrient absorption: enough but not too much. *News In Psychological Sciences* 6, 92–96.
- Diamond, J. (1993) Evolutionary physiology. In: C. A. R. Boyd & D. Noble (eds.) *The Logic of Life*, pp. 89–111. Oxford University Press, Oxford.
- Eastman, J. T. & Coalson, R. E. (1974) The digestive system of the Weddell seal *Leptonychotes weddelli*—a review. In: R. J. Harrison (ed.) *Functional Anatomy of Marine Mammals*. Vol 2, pp. 253–320. Academic Press, London.
- Fadely, B. S., Worthy, G. A. J. & Costa, D. P. (1990) Assimilation efficiency of northern fur seals determined using dietary manganese. *Journal of Wildlife Management* 54, 246–251.
- Fadely, B. S., Zeligs, J. A. & Costa, D. P. (1994) Assimilation efficiencies and maintenance requirements of California sea lions (*Zalophus californianus*) fed Walleye pollock (*Theragra chalcogramma*) and herring (*Clupea harengus*). Final report to National Marine Mammal Laboratory, National Marine Fisheries Service.
- Fisher, K. I., Stewart, R. E. A., Kastelein, R. A. & Campbell, L. D. (1992) Apparent digestive efficiency in walrus (*Odobenus rosmarus*) fed herring (*Clupea harengus*) and clams (*Spisula* sp.). *Canadian Journal of Zoology* 70, 30–36.
- Helm, R. C. (1984) Rate of digestion in three species of pinnipeds. *Canadian Journal of Zoology* 62, 1751–1756.
- Hunt, J. N. & Stubbs, D. F. (1975) The volume and energy content of meals as determinants of gastric emptying. *Journal of Physiology* 245, 209–225.
- Karasov, W. H. & Diamond, J. (1988) Interplay between physiology and ecology in digestion. *Bioscience* 38, 602–611.
- Keiver, K. M., Ronald, K. & Beamish, F. W. H. (1984) Metabolizable energy requirements for maintenance and faecal and urinary losses of juvenile harp seals (*Phoca groenlandica*). *Canadian Journal of Zoology* 62, 769–776.
- Kleiber, M. (1975) *The Fire of Life: An Introduction to Animal Energetics*. Robert E. Krieger Publ. Co., New York.
- Krockenberger, M. B. & Bryden, M. M. (1994) Rate of passage through the alimentary tract of southern elephant seals (*Mirounga leonina*) (Carnivora: Phocidae). *Journal of Zoology, London* 234, 229–237.
- Lavigne, D. M., Barchard, W., Innes, S. & Øritsland, N. A. (1982) Pinniped bioenergetics. In: *Mammals in the Seas*. FAO Fisheries 4(5), 191–235. FAO, Rome.
- Lawson, J. W., Hare, J. A., Noseworthy, E. & Friel, J. K. (1997a) Assimilation efficiency of captive ringed seals (*Phoca hispida*) fed different diets. *Polar Biology* 18, 107–111.
- Lawson, J. W., Miller, E. H. & Noseworthy, E. (1997b) Variation in assimilation efficiency and digestive efficiency of captive harp seals (*Phoca groenlandica*) on different diets. *Canadian Journal of Zoology* 75, 1285–1291.
- Markussen, N. H. (1993) Transit time of digesta in captive harbour seals (*Phoca vitulina*). *Canadian Journal of Zoology* 71, 1071–1073.
- Mårtensson, P.-E., Nordøy, E. S. & Blix, A. S. (1994a) Digestibility of krill (*Euphasia superba* and *Thysanoessa* sp.) in minke whales (*Balaenoptera acutorostrata*) and crabeater seals (*Lobodon carcinophagus*). *British Journal of Nutrition* 72, 713–716.
- Mårtensson, P. E., Nordøy, E. S. & Blix, A. S. (1994b) Digestibility of crustaceans and capelin in harp seals (*Phoca groenlandica*). *Marine Mammal Science* 10, 325–331.

- Martin, R. M. (1977) *Mammals of the Ocean*. G. P. Putnam & Sons, New York.
- Merrick, R. L., Loughlin, T. R. & Calkins, D. G. (1987) Decline in abundance of the northern sea lion, *Eumetopias jubatus*, in Alaska, 1956–1986. *Fisheries Bulletin, United States* **85**, 351–365.
- Miller, L. K. (1978) Energetics of the northern fur seal in relation to climate and food resources of the Bering Sea. *U.S. Marine Mammal Commission, Report No. MMC-75/08*.
- Nordøy, E. S., Sørmo, W. & Blix, A. S. (1993) In vitro digestibility of different prey species of minke whales (*Balenoptera acutorostrata*). *British Journal of Nutrition* **70**, 4485–4489.
- Parsons, J. L. (1977) Metabolic Studies on Ringed Seals (*Phoca hispida*). M.Sc. Thesis, University of Guelph.
- Pitcher, K. W. (1990) Major decline in number of harbor seals, *Phoca vitulina richardsi*, on Tugidak Island, Gulf of Alaska. *Marine Mammal Science* **6**, 121–134.
- Prime, J. H. & Hammond, P. S. (1987) Quantitative assessment of gray seal diet from fecal analysis. In: A. C. Huntley, D. P. Costa, G. A. J. Worthy & M. A. Castellini (eds.) *Approaches to Marine Mammal Energetics*, pp. 165–181. Allen Press, Lawrence.
- Ricker, W. E. (1973) Linear regressions in fishery research. *Journal of the Fisheries Research Board of Canada* **30**, 409–434.
- Ronald, K., Keiver, K. M., Beamish, F. W. H. & Frank, R. (1984) Energy requirements for maintenance and faecal and urinary losses of the grey seal (*Halichoerus grypus*). *Canadian Journal of Zoology* **62**, 1101–1105.
- Rosen, D. A. S. (1996) Seasonal Changes in the Energy Budgets of Captive Harbour Seals (*Phoca vitulina concolor*). Ph.D. Thesis, Memorial University of Newfoundland, St. John's, Nfld.
- Sergeant, D. E. (1973) Feeding, growth, and productivity of Northwest Atlantic harp seals (*Pagophilus groenlandicus*). *Journal of the Fisheries Research Board of Canada* **30**, 17–29.
- Trites, A. W. (1992) Northern fur seals: why have they declined? *Aquatic Mammals* **18**, 3–18.
- Trites, A. W. & Larkin, P. A. (1996) Changes in the abundance of Steller sea lions (*Eumetopias jubatus*) in Alaska from 1956 to 1992: How many were there? *Aquatic Mammals* **22**, 153–166.
- Warner, A. C. I. (1981) Rate of passage of digesta through the gut of mammals and birds. *Nutrition Abstracts and Reviews Series 'B'* **51**, 789–820.